

- Papadopoulos, A. S. T., Igea, J., Smith, T. P. et al. 2019. Ecological speciation in sympatric palms: 4. Demographic analyses support speciation of *Howea* in the face of high gene flow. *Evolution* 73: 1996–2002. doi.org/10.1111/evo.13813
- Sakai, Y. 1964. The species of *Cladophora* from Japan and its vicinity. Scientific papers of the Institute of Algological Research, Faculty of Science, Hokkaido University 5: 1–104
- Seaborn, T. 2014. Limpets and their algal epibionts: costs and benefits of *Acrosiphonia* spp and *Ulva lactuca* growth. *J. Mar. Sci.* 2014: 891943. doi.org/10.1155/2014/891943
- Soejima, A., Yamazaki, N., Nishino, T. & Wakana, I. 2009. Genetic variation and structure of the endangered freshwater benthic alga Marimo, *Aegagropila linnaei* (Ulvophyceae) in Japanese lakes. *Aquat. Ecol.* 43: 359–370. doi.org/10.1007/s10452-008-9204-9
- Suyama, Y., Hirota, S. K., Matsuo, A. et al. 2022. Complementary combination of multiplex high-throughput DNA sequencing for molecular phylogeny. *Ecol. Res.* 37: 171–181. doi.org/10.1111/1440-1703.12270
- Suyama, Y. & Matsuki, Y. 2015. MIG-seq: an effective PCR-based method for genome-wide single-nucleotide polymorphism genotyping using the next-generation sequencing platform. *Sci. Rep.* 5: 16963. doi.org/10.1038/srep16963
- Thines, M. 2019. An evolutionary framework for host shifts-jumping ships for survival. *New Phytol.* 224: 605–617. doi.org/10.1111/nph.16092
- Wahl, M. 1989. Marine epibiosis. I. Fouling and antifouling: some basic aspects. *Mar. Ecol. Prog. Ser.* 58: 175–189. doi.org/10.3354/meps058175
- Wahl, M. 1996. Fouled snails in flow: potential of epibionts on *Littorina littorea* to increase drag and reduce snail growth rates. *Mar. Ecol. Prog. Ser.* 138: 157–168. doi.org/10.3354/meps138157
- Wahl, M. 1997. Increased drag reduces growth of snails: comparison of flume and in situ experiments. *Mar. Ecol. Prog. Ser.* 151: 291–293. doi.org/10.3354/meps151291
- Wahl, M. 2008. Ecological lever and interface ecology: epibiosis modulates the interactions between host and environment. *Biofouling* 24: 427–438. doi.org/10.1080/08927010802339772
- Wahl, M. & Mark, O. 1999. The predominantly facultative nature of epibiosis: experimental and observational evidence. *Mar. Ecol. Prog. Ser.* 187: 59–66. doi.org/10.3354/meps187059
- 王曉陽・阪井与志雄 1986. ツヤナシシオグサとカイゴロモの生活史について. 藻類 34: 209–215
- Ward-Fear, G., Pauly, G. B., Vendetti, J. E. & Shine, R. 2020. Authorship protocols must change to credit citizen scientists. *Trends. Ecol. Evol.* 35: 187–190. doi.org/10.1016/j.tree.2019.10.007
- Williams, G. A. & Morritt, D. 1995. Habitat partitioning and thermal tolerance in a tropical limpet, *Cellana grata*. *Mar. Ecol. Prog. Ser.* 124: 89–103. doi.org/10.3354/meps124089
- Xing, Y. & Wada, K. 2001. Temporal and spatial patterns of the alga *Cladophora conchopheria* on the shell of the intertidal gastropod *Turbo coronatus coreensis*. *Publ. Seto. Mar. Biol. Lab.* 39: 103–111. doi.org/10.5134/176299
- Yamada, M., Wada, K. & Ohno, T. 2003. Observations on the alga *Cladophora conchopheria* on shells of the intertidal gastropod *Turbo coronatus coreensis*. *Benthos. Res.* 58: 1–6. doi.org/10.5179/benthos1996.58.1\_1

(2024年3月6日受付, 2024年5月16日受理)

通信担当編集委員：秋田 晋吾

## 自動会費納入

### 会員各位

会員管理事務の効率化と経費削減のため、自動振替による会費納入にご協力くださいますようお願い申し上げます。当面の間は引き落とし手数料（税別 155 円）を学会の負担とさせていただきます。また、自動振替をご利用の会員全員に領収書を発行させていただいております（送付は 12月末の引き落とし後）。

事務手続き上、外国会員につきましては従来通りクレジットカードによる支払い、また学生会員につきましては郵便振替による振込みのみとさせていただきますので、ご了承ください。振替依頼文書をご希望の方は、下記事務局までご連絡いただきますようお願い申し上げます。

庶務幹事（会員担当）

### 【お申し込み・お問い合わせ先】

〒 729-0292 広島県福山市学園町 1 番地三蔵

福山大学 生命工学部 海洋生物科学科

山岸 幸正 宛

Tel 084-936-2111 内線 4530

E-mail yamagishi@fukuyama-u.ac.jp